

The possibility of a sizable, direct CP-violating asymmetry in $B^\mp \rightarrow K^\mp \eta$

Saul Barshay and Georg Kreyerhoff
 III. Physikalisches Institut
 RWTH Aachen
 D-52056 Aachen
 Germany

February 7, 2008

Abstract

The likelihood of a sizable, direct CP-violating asymmetry in the decays $B^\mp \rightarrow K^\mp \eta$, is calculated within the framework of the model which originally predicted a sizeable asymmetry in $\pi^\mp \eta(\eta')$. It is shown in a transparent manner that these decays are the best places for providing the first clear evidence for direct CP violation in the decays of a charged particle. These decays contain two amplitudes of comparable magnitude, with different weak-interaction phase, and with different strong-interaction phase which is explicitly calculated in a model of three coupled channels of physical hadrons.

Recently, the BABAR collaboration reported the observation of the decays $B^\mp \rightarrow \pi^\mp \eta$, with a small branching ratio of $(4.2 \pm 1.0 \pm 0.3) \times 10^{-6}$ [1]. It is notable that there is an indication of a sizable, direct CP-violating asymmetry $\mathcal{A}(B^- - B^+) = (-0.51 \pm 0.19 \pm 0.01)$ [1]. At present, this is potentially the first distinctly nonzero asymmetry ever detected in the decay of a charged particle. That this particular decay mode with a small branching ratio provides the best possibility for observing direct CP violation in charged-meson decays, was predicted in 1991 [2]. The prediction is based upon detailed calculations of partial rates and asymmetries, which utilize known, strongly-coupled channels of physical hadrons $\pi^\mp(\eta, \eta', \eta_c)$ in order to calculate the final-state interaction phases which are essential for obtaining a direct, CP-violating asymmetry [2, 3]. The other necessary ingredient [2, 3] is the presence of two weak, effective amplitudes, each with a different CP-violating phase (given as functions of the phase δ_{13} of the CKM quark-mixing matrix). The CP-violating asymmetry arises from the interference between these two amplitudes; it can be sizable when the amplitudes have comparable magnitude [2, 3]. This is the case for $B^\mp \rightarrow \pi^\mp \eta(\eta')$, because one tree-level diagram ^{F1} for $b \rightarrow u + (W \rightarrow \bar{u}d)$ is proportional to $|s_{13}e^{-i\delta_{13}}| \sim |(s_{12}s_{23}/2.5)e^{-i\delta_{13}}|$; and the second tree-level diagram for $b \rightarrow c + (W \rightarrow \bar{c}d)$ is proportional to $|s_{23}s_{12}|$. From the strong-interaction transition $\eta_c \pi \rightarrow \eta(\eta')\pi$, the second tree diagram acquires a multiplication factor of $+i$ times an effective phase angle, approximately calculated [2] to be a number of the order of 0.03.^{F2} From interference with the first tree diagram, one immediately obtains an order of magnitude estimate for the direct, CP-violating asymmetry,

^{F1}The s_{ij} denote sines of quark-mixing angles in the usual form [4] of the CKM matrix, with δ_{13} the large, CP-violating phase; $\alpha_s \sim 0.2$ is the QCD coupling at a scale $\sim m_{B^0} \sim 5$ GeV [4]. Numerical values of the s_{ij} and δ_{13} are taken from [4].

^{F2}In the original paper [2], the K -matrix elements are underestimated by a factor of 2. Incorporating this factor, increases the two direct, CP-violating asymmetries for η, η' by a factor of 2. (The CP-violating phase denoted by ϕ is δ_{13} .)

$\mathcal{A} \sim -\frac{2(0.03)(1/2.5)}{(1/2.5)^2} \sin \delta_{13} \sim -0.15 \sin \delta_{13}$. For $B^\mp \rightarrow \pi^\mp \eta(\eta')$, the loop diagrams involving $b \rightarrow t, c$ and $t, c \rightarrow d$, with gluon emission and conversion into a $\bar{u}u$ quark pair are proportional to $|s_{12}s_{23}| \times \alpha_s/\pi$, and so are down by a factor of the order of $\alpha_s/\pi \sim 0.07$. The small branching ratio for $B^\mp \rightarrow \pi^\mp \eta'$ has very recently been reported [5], $(2.8_{-1.0}^{+1.3} \pm 0.6) \times 10^{-6}$. It appears to be comparable to, but a little less than the branching ratio for $\pi^\mp \eta$, as predicted [2]. A sizable CP-violating asymmetry is expected [2].

In this note, we discuss three positive and one negative result which follow directly from the model which led to the above predictions, when the model is extended to the decays $B^\mp \rightarrow K^\mp(\eta, \eta', \eta_c)$. Within the context of the quark diagrams considered in the model, we discuss two possible remedies for the negative result which concerns the quite large empirical branching ratio for the decays $B^\mp \rightarrow K^\mp \eta'$. We then consider the explanation in terms of additional quark diagrams, namely the weak loop (“penguin”) diagrams, involving $b \rightarrow t, c$ and $t, c \rightarrow s$, with gluon emission and conversion into an $\bar{s}s, \bar{u}u$ pair. Such diagrams are generally considered to have an essential role [6] in the decays $B \rightarrow \pi K$, and also in $B^\mp \rightarrow K^\mp \eta(\eta')$ [7]. Explanations of the large η' branching ratio in the model without additional loop diagrams are possible, but do not seem to be adequate. Contributions from both dynamics are probably necessary. A new contribution involving a small $\bar{c}c$ component in η' is considered, and can be tested for experimentally [3]. Our main positive results concern the empirically small branching ratio for $B^\mp \rightarrow K^\mp \eta$, and most importantly for the experimental situation [1], the possibility of a sizable, direct CP-violating asymmetry $\mathcal{A} \sim -0.2 \sin \delta_{13}$, for this decay mode. The recent report [1] which indicated a sizable asymmetry for $\pi^\mp \eta$, also gave an asymmetry for $K^\mp \eta$, of $-0.32 \pm 0.20 \pm 0.01$. This is a principal motivation for the present analysis, because this could be a second place where sizeable, direct CP violation can be anticipated and observed in the decay of a charged meson. It is valuable for experimentalists to have a transparent estimate for direct CP violation in charged-meson decay for those few modes where a sizable asymmetry can likely be clearly observed [1]. Such clear observations would be a milestone in the history of CP noninvariance [8, 9, 10].

The first result follows immediately from the number obtained in 1991 for the branching ratio for $B^\mp \rightarrow \pi^\mp \eta_c$, given as about 4.1×10^{-5} for the decay constant $f_{\eta_c} \sim 310$ MeV. From this number, the branching ratio for $B^\mp \rightarrow K^\mp \eta_c$ is obtained approximately by simply dividing by $s_{12}^2 \cong 0.48$; it is thus predicted to be quite large, $\sim 1 \times 10^{-3}$ (for $f_{\eta_c} \cong 335$ MeV [11]). Recent experimental results confirm this: CLEO [11] gives $(0.69_{-0.21}^{+0.26} \pm 0.08 \pm 0.20) \times 10^{-3}$; BELLE [12] gives $(1.25 \pm 0.14_{-0.12}^{+0.10} \pm 0.38) \times 10^{-3}$. The first error is statistical, the second is systematic, and the third is from the η_c branching-fraction uncertainty. The large branching ratio supports the essential validity of the phenomenological, effective weak hamiltonian [13], and of the physical hadron “factorization” approximation [13, 14, 15] used in calculating matrix elements for particular exclusive decay modes (that is, matrix elements of products of weak currents factorize into matrix elements of individual currents in states of physical hadrons). We explain next the second and third results, concerning the small branching ratio for $B^\mp \rightarrow K^\mp \eta$, and concerning the distinct possibility of a sizable, direct CP-violating asymmetry. From eq. (27) in [2], the form of the full, “unitarized” [16] amplitude (denoted by A^u) ^{F3} is ^{F2}

$$\begin{aligned} A_\eta^u &\cong A_\eta + it_{12}A_{\eta'} + it_{13}A_{\eta_c} \\ &\sim A_\eta - i(0.026)A_{\eta_c} \end{aligned} \quad (1)$$

The A are the weak amplitudes following from the effective, weak hamiltonian, com-

^{F3}The three amplitudes $A_{\eta, \eta', \eta_c}^u$ satisfy CPT invariance [2], namely that the widths of B^- and B^+ , summed over all three channels $K^\mp(\eta, \eta', \eta_c)$ are equal.

puted with the factorization approximation. The t_{ij} ($i, j = 1, 2, 3 = \eta, \eta', \eta_c$) are given approximately [2] by linear combinations of the tangents of the three small eigenphases; these were originally calculated [2] for the specific system of three coupled channels $\pi^\mp(\eta, \eta', \eta_c)$. The essential number is^{F2} $|t_{13}| \sim 0.026$ calculated in [2]; we assume that this has about the same magnitude for the system of three coupled channels, $K^\mp(\eta, \eta', \eta_c)$.^{F4} There is a sign ambiguity related to the analogue for $K^\mp$ of eq. (18) in [2] for $\pi^\mp$. Here we choose the negative sign, which leads to a negative t_{13} . Except for the CKM factors, the magnitude of the weak amplitudes are taken as approximately the same as calculated in [2].^{F5} In eq. (1), A_{η_c} is proportional to $|s_{12}s_{13}e^{-i\delta_{13}}| \sim \left|\frac{s_{12}^2 s_{23}}{2.5}e^{-i\delta_{13}}\right| \sim (0.02)s_{23}$. In this initial discussion, we neglect the term in $A_{\eta'}$ which, in the present approximation, is similar in magnitude [2] to A_η , and is multiplied by the calculated [2] small number $|t_{12}| \sim 0.1$.^{F2} Below, we discuss the addition of the loop diagram that can give a large $A_{\eta'}$ in eq. (1), and which can then make this term comparable to the first term (which is also enhanced somewhat). Our main result here, concerning the possibility of a sizable CP-violating asymmetry in $B^\mp \rightarrow K^\mp \eta$, will not be changed by inclusion of an enhanced $A_{\eta'}$. The branching ratio for $K^\mp \eta$ is increased, and the asymmetry correspondingly reduced. Computing the explicit numerical value [2] of A_η , and expressing it as a multiple of the product of the amplitude arising from A_{η_c} and the strong final-state interaction factor, we obtain^{F2}

$$A_\eta^u \sim - \left\{ \left(\frac{17}{39} \right) e^{-i\delta_{13}} + i \right\} (0.026) A_{\eta_c} \quad (2)$$

Thus, there is a clear demonstration of two amplitudes of comparable magnitude, differing in weak-interaction phase and in strong-interaction phase. With the large branching ratio for $K^\mp \eta_c$ of about 10^{-3} , this gives the small branching ratio for $K^\mp \eta$ of about

$$\begin{aligned} & \frac{1}{2} \{ \text{b.r.}(B^- \rightarrow K^- \eta) + \text{b.r.}(B^+ \rightarrow K^+ \eta) \} \\ & \sim \{ 1 + (17/39)^2 \} \times (0.026)^2 \times 10^{-3} \cong 0.8 \times 10^{-6} \end{aligned} \quad (3)$$

The model gives a quite small branching ratio for $K^\mp \eta$, essentially in terms of the large branching ratio for $K^\mp \eta_c$ and a calculated, small final-state interaction factor. Recent measurement [1] gives for the branching ratio for $K^\mp \eta$, $(2.8 \pm 0.8 \pm 0.2) \times 10^{-6}$. From eq. (2) the CP-violating asymmetry is calculated as

$$\mathcal{A}(B^- - B^+) \cong - \frac{2(17/39) \sin \delta_{13}}{\{1 + (17/39)^2\}} \sim -0.73 \sin \delta_{13} \quad (4)$$

The possibility of an asymmetry which is quite sizable in magnitude occurs naturally in this model.

The negative result within the model is a small branching ratio for $B^\mp \rightarrow K^\mp \eta'$. The analogue of eqs. (1,2) is [2]^{F2}

$$\begin{aligned} A_{\eta'}^u & \cong A_{\eta'} + it_{21} A_\eta + it_{23} A_{\eta_c} \cong A_{\eta'} - i(0.04) A_{\eta_c} \\ & \sim - \left\{ \left(\frac{8}{39} \right) e^{-i\delta_{13}} + i \right\} (0.04) A_{\eta_c} \\ & \Rightarrow \frac{1}{2} \{ \text{b.r.}(B^- \rightarrow K^- \eta') + \text{b.r.}(B^+ \rightarrow K^+ \eta') \} \sim 1.7 \times 10^{-6} \end{aligned} \quad (5)$$

^{F4}The experimental limit [4] for the branching ratio for $\eta_c \rightarrow K \bar{K} \eta$ is $< 3.1\%$, comparable to the measured $(4.9 \pm 1.8)\%$ for $\pi \pi \eta$ and $(4.1 \pm 1.7)\%$ for $\pi \pi \eta'$. (That for $K \bar{K} \pi$ is $(5.5 \pm 1.7)\%$.) The t_{13}, t_{23} are related to the square root of the branching ratios [2]. Only if $\eta_c \rightarrow K \bar{K} \eta(\eta')$ were to be very small would they be much smaller than used here, and hence the asymmetry in $K^\mp \eta$ smaller.

^{F5}Using the approximation of the same decay constant and wave-function overlap for $K^\mp$ as for $\pi^\mp$, as given in [2] ($f_{\pi^+} \sim 130$ MeV [4]).

The experimental numbers are about 40 times larger, $(6.5 \pm 1.5 \pm 0.9) \times 10^{-5}$ and $(80 \pm 10 \pm 79 \times 10^{-6})$ from CLEO [17], $(79 \pm 12 \pm 9) \times 10^{-6}$ from BELLE [18], and $(70 \pm 8 \pm 5) \times 10^{-6}$ from BABAR [19]. So a factor of about 6.3 larger in amplitude. Within the context of the quark diagrams considered so far in the model, there are two ways to increase the amplitude.

- (1) Constructive contributions from additional (to η_c) $\bar{c}c$ states could contribute to enhancing an effective $|t_{23}|$. Specifically, the state χ_{c1} and the higher-mass η_c state(s) may have significant decay amplitudes to $\eta' K^+ K^-$. It is notable that χ_{c1} does have significant, measured [4] branching ratios to $\pi^- K^+ K_S^0$ and to $\pi^+ \pi^- K^+ K^-$. However, invoking a hypothetical factor of 3 increase in an effective $|t_{23}|$, to ~ 0.12 , still leaves the branching ratio of $\sim 14 \times 10^{-6}$ below the experimental value.
- (2) There could be a small $\bar{c}c$ component in η' . Applied to η , this is an old idea due to Feldman and Perl [20], who considered it as a possible explanation for the relatively large branching ratio for $\Psi' \rightarrow J/\Psi \eta$. Such a component, at the level of $< 15\%$ in amplitude, was used in [3] to calculate two weak amplitudes of comparable magnitude, whose interference led to sizable, direct CP-violating asymmetries in the angular distributions and rates for $B^0(\bar{B}^0) \rightarrow \pi^+ \pi^- \eta$. However, invoking a hypothetical $\bar{c}c$ amplitude in η' of about 15%, again raises the $K^\mp \eta'$ branching ratio to only about 23×10^{-6} . A direct experimental test for such an amplitude would be in the then expected [3] sizable, direct CP violation in angular distributions and rates for $B^0(\bar{B}^0) \rightarrow \pi^- \pi^+ \eta'$.

We turn to the weak loop diagrams involving $b \rightarrow t, c$ and $t, c \rightarrow s$, which are proportional to $s_{23}(\alpha_s/\pi)$ and which seem to be necessary to explain the large branching ratio for $K^\mp \eta'$. Our main concern here is the implication of an enhanced η' amplitude for the direct, CP-violating asymmetry estimated above in eq. (4) for the small-branching mode $K^\mp \eta$. Lipkin noted [7] that the negative sign of the $\bar{s}s$ amplitude in the η , as compared to the positive sign in the η' , simply leads to a strong suppression of $B^\mp \rightarrow K^\mp \eta$, when the decay amplitude involves the $b \rightarrow s + (g \rightarrow \bar{s}s)$ loop diagram with gluon emission and conversion into an $\bar{s}s$ pair. Indeed, if this decay amplitude dominates and is responsible for an experimental [17, 18, 19] b.r. $(K^\mp \eta') \sim 70 \times 10^{-6}$, then a possible suppression factor [7] of the order of $1/34$ brings one to $1/2\{\text{b.r.}(K^- \eta) + \text{b.r.}(K^+ \eta)\}$ of $\sim 2 \times 10^{-6}$, close to the experimental [1] number. This squared amplitude would increase the numerical value of the denominator in eq. (4) by a factor of the order of 3. If this amplitude contributes little to the numerator (i. e. if it has negligible phase factors), then the asymmetry in eq. (4) would be reduced to $\mathcal{A} \sim -0.24 \sin \delta_{13}$. With an enhanced weak amplitude $A_{\eta'}$, the term $it_{12}A_{\eta'}$ in eq. (1) also contributes to A_η^u (with $|t_{12}| \sim 0.1)^{F6}$. This term could increase the numerator in eq. (4), as well as the denominator; including it constructively, we estimate $\mathcal{A} \sim -0.2 \sin \delta_{13}$. It is an open question [17] as to whether the loop diagrams alone can actually account for the η' branching ratio. Assuming $b \rightarrow s + g$ to have a branching fraction of about 0.5% ^{F6}, then a (quasi) exclusive mode, like $(K^\mp \pi^0 + K^0 \pi^\mp)$, if largely “penguin” mediated, would have a branching ratio of the order of $(1/10)(\alpha_s/\pi)(5 \times 10^{-3}) \sim 3.5 \times 10^{-5}$, with $\sim \alpha_s/\pi \sim 0.07$ arising from the gluon emission and conversion into $\bar{u}u, \bar{d}d$, and the factor of $\sim 1/10$ approximately accounting for a (quasi)exclusive mode. ^{F6} This simple estimate is remarkably close to experiment [4] for $K\pi$, but is still somewhat below $K^\mp \eta'$ [16]. In addition, there is a small contribution from the amplitudes

^{F6}Note that reducing a branching ratio of $\sim 5 \times 10^{-3}$ by $\alpha/\alpha_s \sim (1/137)/0.2 \cong 1/27$ results in $\sim 1.8 \times 10^{-4}$, only just below the experimental branching ratio [4] of $(3.1 \pm 1.1) \times 10^{-4}$ for $b \rightarrow s + \gamma$. A dominant exclusive channel $K^*(892)\gamma$, is down by $1/8$ experimentally [4].

giving rise to eq. (5). In this situation, a diagram involving a hypothetical, small (up to $\sim 18\%$ in amplitude) $\bar{c}c$ component [20, 3] in η' , if constructive with the loop diagram, could increase the branching ratio to experiment. As mentioned above, a direct experimental test for such a component is possible in the decays $B^0(\bar{B}^0) \rightarrow \pi^-\pi^+\eta'$. The effect upon the $K^\mp\eta$ asymmetry is to increase it somewhat (~ 1.4), because of a decrease in the calculated $K^\mp\eta$ partial rate from the smaller loop contribution. In contrast to the prediction of a sizable, direct CP-violating asymmetry in $K^\mp\eta$, the prediction for $K^\mp\eta'$ is for a small asymmetry, of the order of a few percent. This is because of the dominant size of the loop diagram (approximately independent of δ_{13}), in determining the branching ratio. An interference term normalized to the partial rates, giving rise to an asymmetry, must be down by a factor of at least of the order of 1/10 from eq. (4). Interestingly, the experimental number [4], appears to be small, $\mathcal{A}_{K^\mp\eta'} = -0.02 \pm 0.07$.

In summary, in addition to the original prediction [2] for $B^\mp \rightarrow \pi^\mp\eta(\eta')$, the extended model predicts the likelihood of a sizable, direct CP-violating asymmetry in $B^\mp \rightarrow K^\mp\eta$, but not in $K^\mp\eta'$. The physical origin of these predictions is the presence in the three decay processes of two amplitudes which are comparable in magnitude, and which contain different CP-violating weak phases and different strong-interaction phases [3]. The small comparable magnitudes result from small relevant factors in the CKM matrix and from the small magnitude of (imaginary) numbers which are obtained from calculating [2] the strong-interaction phases in a specific model of three coupled channels of physical hadrons.^{F7} Decay modes with relatively large branching ratios, like $K^\mp\eta'$, tend to be dominated by a single, large decay amplitude, and hence have small asymmetries. Thus, $B^\mp \rightarrow \pi^\mp\eta(\eta')$ and likely^{F4} $K^\mp\eta$, are the best processes in which direct CP violation can be established for the first time in the decay of a charged particle.

S. B. thanks Lahlit Sehgal for information and constructive criticism.

References

- [1] BABAR Collab., B. Aubert et al., hep-ex/0303039
- [2] S. Barshay, D. Rein and L. M. Sehgal, Phys. Lett. **B259** (1991) 475
- [3] S. Barshay and J. Geris, Phys. Lett. **B84** (1979) 150
- [4] Particle Data Group Collab., K. Hagiwara et al., Phys. Rev. **D66** (2002) 01001-1
- [5] BABAR Collab., B. Aubert et al., hep-ex/0308015
- [6] N. G. Deshpande et al., Phys. Rev. Lett. **82** (1999) 2240
- [7] H. J. Lipkin, Phys. Lett. **B254** (1991) 247
- [8] J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turlay, Phys. Rev. Lett. **13** (1964) 138
- [9] NA31 Collab., H. Burkhardt et al., Phys. Lett. **B206** (1988) 168
NA48 Collab., V. Fanti et al., Phys. Lett. **B465** (1999) 335
- [10] KTeV Collab., A. Alevi-Horati et al., Phys. Rev. Lett. **38** (1999) 22

^{F7}The use of coupled channels for calculating strong-interaction phases has been applied in trying to understand recent data on CP violation in $B^0(\bar{B}^0) \rightarrow \pi^+\pi^-$. S. Barshay and G. Kreyerhoff, hep-ph/0303254v4; JHEP in press.

- [11] CLEO Collab. K. W. Edwards et al., Phys. Rev. Lett. **86** (2001) 30
- [12] BELLE Collab., F. Fang et al., Phys. Rev. Lett. **90** (2003) 071801-1
- [13] M. Bauer, B. Stech and M. Wirbel, Z. Phys. **C34** (1987) 103
- [14] S. Barshay and L. M. Sehgal, Phys. Lett. **B185** (1987) 150
- [15] S. Barshay, E. Eich and L. M. Sehgal, Phys. Lett. **B202** (1988) 202; **B212** (1988) 542 (E)
- [16] A. N. Kamal, N. Sinha and R. Sinha, Z. Phys. **C41** (1988) 207
- [17] CLEO Collab., B. H. Behrens et al., Phys. Rev. Lett. **80** (1998) 3710
CLEO Collab., S. J. Richichi et al., Phys. Rev. Lett. **85** (2000) 520. These papers refer to many theoretical papers. Estimates range from about 7 to 41×10^{-6} .
- [18] BELLE Collab., K. Abe et al., Phys. Lett. B517 (2001) 309
- [19] BABAR Collab., B. Aubert et al., hep-ex/0108017v2
- [20] G. L. Feldmann and M. L. Perl, Phys. Rep. **33C** (1977) 285n